



Rochester (248) 650-2400 Shelby Township (586) 254-2777 Farmington Hills (248) 865-4238 River District (810) 329-1250

# Post Surgery Instructions Minimally Invasive Hallux Valqus (Bunion) Correction

#### **General Treatment Facts**

- The goal of this surgery is to reduce pain and improve the alignment and function of the great toe joint.
- During this procedure, the metatarsal (toe) bone is cut and shifted in order to straighten the toe. The big toe ligaments are loosened on one side and tightened on the other to straighten the toe.
- You will be able to walk on the heel and outside of the foot the day of surgery while wearing a special postoperative shoe.
- You will experience significant swelling in the foot after surgery. This can last up to 4 to 6 months. The more that you elevate the foot and limit your walking in the first 2 weeks, the better your pain and swelling will be.
- The post-operative shoe must be worn for 6-12 weeks, depending on your specific procedure.
- You will not be able to drive until 6 weeks if it is your right foot.

### **Post Surgery Course**

## Day 1

- A sterile dressing will be applied and the foot will be placed in a protective shoe.
- Please do not remove the dressing. If for any reason the dressing is too tight, loosen it and re-apply it to your comfort level.
- It is important to ice and elevate the foot, take pain medication and rest as needed.
- Do not get the toe or dressing wet.
- Move the ankle, knee, hip, and upper extremities as tolerated.

#### Day 4

• Pain should be better after the 3<sup>rd</sup> day. If your pain worsens since day 3 or you have a fever and/or chills, please call the office.

# 2 Weeks (approximately)

- First post-operative visit.
- Your dressing will be taken off and your sutures will be removed.
- Continue to walk on the heel and outside of the foot only in the postop shoe.
- Your activity level will be about 30% of your normal activity level.

## 3 Weeks

- Gentle exercise on a stationary bike is permitted.
- Continue to wear the post-operative shoe and do not roll through the front of the foot.

#### 6 Weeks

- Wean out of the protective shoe.
- It is normal to experience foot swelling at and beyond this point. You may require a shoe that is one to two sizes larger than your normal shoe size. Otherwise, you may need to loosen your shoelaces to accommodate the swelling.
- You may drive in a supportive tennis shoe.
- If you have pins in your toes, they are usually pulled at this visit. You can get the foot wet 5-7 days after your pins are pulled.
- Avoid walking on uneven or rocky terrain.
- You can exercise on a bike as long as it's on the arch of the foot.

# 12 Weeks

- Ambulate as tolerated in a comfortable, supportive tennis shoe.
- Increase activity as tolerated.